

High-order corrections to the holographic entropy bound from quantum field theory

Yi Ling*

*Center for Gravity and Relativistic Astrophysics,
Department of Physics, Nanchang University, Nanchang, 330047, China
CCAST (World Laboratory), P.O. Box 8730, Beijing, 100080, China*

Hongbao Zhang†

*Department of Astronomy, Beijing Normal University, Beijing, 100875, China
Department of Physics, Beijing Normal University, Beijing, 100875, China
CCAST (World Laboratory), P.O. Box 8730, Beijing, 100080, China*

(Dated: February 7, 2008)

From the viewpoint of local quantum field theory, this letter investigates the high-order corrections to the holographic entropy bound. As a result, the logarithmic correction term appears naturally with the definite coefficient $-\frac{1}{2}$, which thus provides another semi-classical platform to examine such candidates for quantum gravity as Loop Quantum Gravity and String Theory.

PACS numbers: 04.70.-s, 04.62.+v, 04.60.-m, 03.67.-a

In a recent paper[1], Yurtsever provides an explanation for the holographic principle from local quantum field theory. Based on the cutoff of the one-mode's energy by the Planck scale physics and the constraint of the total energy of Fock states against gravitational collapse, Yurtsever shows that the leading term in the entropy-area relation is linear, which is in full agreement with the well established holographic entropy bound. However, in recent years evidence has been mounting that smaller correction terms may also exist. The purpose of this letter is to further investigate the high-order corrections to the holographic entropy bound from the viewpoint of local quantum field theory.

To proceed, we shall confine ourselves to the real massless scalar in the spherical region with the radius R , which is probably the more natural choice than the cube considered by Yurtsever. In addition, as is done by Yurtsever, we assume that the maximum energy of admissible one-modes is of the order of the Planck energy, i.e. given by Λ , where Λ is of the order of one for naive Planck scale physics. Moreover, the maximum energy of a state in the symmetric Fock space should be constrained by the energy of a Schwarzschild black hole with radius R , i.e. $\frac{R}{2}$.

Then by the simplified derivation of Yurtsever's result[2], the dimension of the gravitational truncated symmetric Fock space approximately reads

$$W = I_0(2\sqrt{z}). \quad (1)$$

Here I_0 is the zeroth-order Bessel function of the second kind, and z is obtained by

$$z = \int_0^\Lambda d\omega \rho(\omega) \frac{R}{2\omega} = \frac{\Lambda^2}{6\pi \cdot (4\pi)^2} A^2, \quad (2)$$

where the density of states $\rho(\omega) = \frac{V}{2\pi^2} \omega^2$ has been employed with the volume $V = \frac{4\pi}{3} R^3$ and the area $A = 4\pi R^2$. Later, according to the asymptotic behavior of I_0 as $x \rightarrow \infty$, i.e.,

$$I_0(x) \sim \frac{e^x}{\sqrt{2\pi x}}, \quad (3)$$

the maximum entropy takes the form

$$S_{max} = \ln W \sim \frac{2\Lambda}{4\pi \cdot \sqrt{6\pi}} A - \frac{1}{2} \ln A. \quad (4)$$

Some remarks on this result are presented in order. Firstly the leading term is obviously dependent on the the Planck scale physics where the ordinary quantum field theory is broken. Specifically, if $\Lambda = \frac{\pi \cdot \sqrt{6\pi}}{2}$, the leading term is the just the famous linear entropy-area relation formula. While the sub-leading correction is independent of the Planck scale physics. In particular, it is worth noting that the logarithmic term has also arisen in both Loop Quantum Gravity and String Theory. However, one should emphasize that there is no general agreement on the coefficient of the logarithmic correction[3]. Therefore the resultant coefficient $-\frac{1}{2}$ obtained here acquires much importance: since our foregoing assumptions only rely on classical general relativity and quantum field theory below the Planck scale, it provides another semi-classical constraint onto all candidates for quantum gravity such as Loop Quantum Gravity and String Theory.

In summary, based on the viewpoint of local quantum field theory, which was initially proposed by Yurtsever to explain the famous linear entropy-area relation, we have shown that the logarithmic correction to the holographic entropy naturally appears with the definite coefficient $-\frac{1}{2}$, which should serve as another severe semi-classical constraint onto any candidate for quantum gravity.

*Electronic address: yling@ncu.edu.cn

†Electronic address: hzbzhang@pku.edu.cn

Acknowledgements

Y. Ling's work is partly supported by NSFC (Nos.10405027 and 10205002) and SRF for ROCS,

SEM. H. Zhang's work is supported in part by NSFC(Nos.10373003 and 10533010).

-
- [1] U. Yurtsever, Phys. Rev. Lett. 91: 041302(2003).
[2] A. Aste, Europhys. Lett. 69: 36-40(2005).

- [3] S. Hod, Class. Quant. Grav. 21: L97-L100(2004).